

Table 22. Reported cases of HIV infection (not AIDS) for female adults and adolescents, by transmission category and race/ethnicity, cumulative through 2004—42 areas with confidential name-based HIV infection reporting

Transmission category	White, not Hispanic				Black, not Hispanic				Hispanic			
	2004		Cumulative through 2004 ^a		2004		Cumulative through 2004 ^a		2004		Cumulative through 2004 ^a	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Injection drug use	341	21	3,513	26	546	9	6,223	14	368	19	1,816	19
Hemophilia/coagulation disorder	0	0	20	0	5	0	31	0	2	0	9	0
Heterosexual contact	667	40	6,236	46	2,358	39	19,445	45	1,003	51	4,505	48
<i>Sex with injection drug user</i>	147	9	1,685	12	273	4	3,269	8	233	12	1,018	11
<i>Sex with bisexual male</i>	50	3	540	4	110	2	1,075	2	27	1	177	2
<i>Sex with person with hemophilia</i>	5	0	106	1	8	0	63	0	1	0	9	0
<i>Sex with HIV-infected transfusion recipient</i>	5	0	53	0	19	0	100	0	2	0	31	0
<i>Sex with HIV-infected person, risk factor not specified</i>	460	28	3,852	28	1,948	32	14,938	35	740	38	3,270	35
Receipt of blood transfusion, blood components, or tissue	9	1	144	1	43	1	332	1	12	1	62	1
Other/risk factor not reported or identified	646	39	3,772	28	3,144	52	17,254	40	576	29	3,029	32
Total	1,663	100	13,685	100	6,096	100	43,285	100	1,961	100	9,421	100

Transmission category	Asian/Pacific Islander				American Indian/Alaska Native				Total			
	2004		Cumulative through 2004 ^a		2004		Cumulative through 2004 ^a		2004 ^b		Cumulative through 2004 ^a	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Injection drug use	3	4	18	5	3	8	87	28	1,267	13	11,719	17
Hemophilia/coagulation disorder	0	0	1	0	0	0	0	0	7	0	62	0
Heterosexual contact	30	40	138	42	17	45	139	45	4,090	41	30,644	45
<i>Sex with injection drug user</i>	3	4	10	3	2	5	49	16	666	7	6,063	9
<i>Sex with bisexual male</i>	1	1	7	2	1	3	10	3	189	2	1,822	3
<i>Sex with person with hemophilia</i>	0	0	0	0	0	0	2	1	14	0	182	0
<i>Sex with HIV-infected transfusion recipient</i>	0	0	1	0	0	0	0	0	26	0	186	0
<i>Sex with HIV-infected person, risk factor not specified</i>	26	35	120	36	14	37	78	25	3,195	32	22,391	33
Receipt of blood transfusion, blood components, or tissue	0	0	5	2	0	0	1	0	64	1	546	1
Other/risk factor not reported or identified	42	56	169	51	18	47	80	26	4,446	45	24,572	36
Total	75	100	331	100	38	100	307	100	9,874	100	67,543	100

Note. Includes only persons with HIV infection that has not progressed to AIDS.

Since 2004, the following 42 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Connecticut, Florida, Georgia, Idaho, Indiana, Iowa, Kansas, Kentucky, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, Puerto Rico, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, American Samoa, Guam, Northern Mariana Islands, and the U.S. Virgin Islands.

Connecticut has confidential name-based HIV infection reporting only for pediatric cases.

Florida (since July 1997) has had confidential name-based HIV infection reporting only for new diagnoses.

Pennsylvania (October 2002) implemented confidential name-based HIV infection reporting only in areas outside the city of Philadelphia.

Texas (February 1994 through December 1998) reported only pediatric HIV infection cases.

^a Includes persons with a diagnosis of HIV infection (not AIDS), reported from the beginning of the epidemic through December 2004. Cumulative total includes 514 females of unknown race or multiple races.

^b Total includes 41 females of unknown race or multiple races.